

Product description

A novel syngeneic Pik3ca+.p53+/- murine lung cancer cell line (Y856) with orthotopic growth potential, that can serve as a useful preclinical model for evaluating tumour-host interactions, the impact of specific oncogenic alterations on the tumour microenvironment, and immunotherapeutic approaches. This cell line was derived from the primary lung tumour of a R26StopFLP110*fl/+.p53fl/+ female C57BL/6 mouse and can form orthotopic tumours in immunocompetent C57BL/6 mice. Y856 cells demonstrate genetic recombination at the Pik3ca and Tp53 loci as shown by PCR and show reduced expression of TP53 and consistent with constitutive PIK3CA activation, show increased pAKT expression without increased total AKT, as shown by western analysis.

Name: Pik3ca+.p53+/- murine lung cancer cell line

Organism: Mouse

Gender: Female

Tissue: Lung

Disease: Cancer

Cancer: Lung cancer

Growth properties: Adherent

Production details: A 20-week-old R26StopFLP110* fl/+.p53fl/+ female mouse was injected with 2 µl of 10⁹ PFU/ml Ad5-CMV-Cre into the left lung. When this animal was euthanized 11 week later, a 6 mm left lung tumour was combined with multiple metastasis (contralateral lung, pericardial, pleural) then passaged into a female recipient animal. After one passage through the flank, the Pik3ca+.p53+/- cell line (Y856) was established; this line could form orthotopic tumours in immunocompetent C57BL/6 animals.

Application: 2D cell culture; Immunohistochemistry (IHC); Cell line-derived xenograft (CDX) model development

Biosafety level: 1

Contributor(s)

Inventor: Stephen Malkoski

Institute: The University of Colorado Anschutz Medical Campus

Properties

Product format: Frozen

Unpacking and storage:

1. Check all containers for leakage or breakage.
2. Remove the frozen cells from the dry ice packaging and immediately place the cells at a temperature below -130°C, preferably in liquid nitrogen vapor, until ready for use.

Recommended medium: Dulbecco's Modified Eagle Medium (DMEM) containing 4.5 g/L d-glucose, L-glutamine, and sodium pyruvate, supplemented with 10% (v/v) fetal bovine serum and 100 µg/mL of primocin

(optional).

Subculture: 0.25% Trypsin-EDTA at RT for 2-10 minutes or 37°C if especially hard to detach. Split 1:5 to 1:15 when 90% confluent.

Culture conditions: 37.0°C ± 1.0°C humidified incubator with 5.0% ± 1.0% CO₂.

Handling instructions

1. Please ensure that vials are frozen when received, and store at **<-130 °C long term**. When removing frozen cells from storage, it is important to minimize exposure to room temperature (15 - 25°C). If not proceeding directly to thawing, place the cells on dry ice or in a liquid nitrogen container.
2. **Do not thaw at room temperature.** To thaw, swirl the vial quickly in a 37 °C water bath with O-ring and cap above the water to avoid contamination. Remove from the water bath with a small ice pellet remaining (this should not take more than 2 minutes) and wipe the exterior with 70% ethanol or isopropanol before transferring to a biosafety cabinet. Further steps should be conducted under aseptic conditions.
3. We strongly recommend that the volume of cell suspension is measured at this point, and a 20 uL aliquot be removed for a **viable cell count** using trypan blue or similar dye. This ensures that provided cells are viable, and the cell count can be used to determine volume of growth medium to be added to the cell suspension.
4. Transfer the cell suspension into a 15 mL sterile conical tube containing 5 mL of pre-warmed, complete growth medium. Centrifuge cells at **125xg for 5-7 minutes**.
5. Aspirate the supernatant without disturbing the cell pellet. Re-suspend the cell pellet in the recommended pre-warmed, complete growth medium and dispense into a T25 culture flask.
6. Seed at ~20,000 to 50,000 cells/cm² (seed more densely at initial thaw - cells can tolerate more sparse subcultures once proliferating consistently) incubate the cells at the recommended conditions.

References

- Nolan et al. 2020. Cancer Cell Int. 28: 20:417. PMID: 32874131

Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner:
Pik3ca+.p53+/- murine lung cancer cell line, was invented by Stephen Malkoski at The University of Colorado Anschutz Medical Campus (CancerTools.org #162388).

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

While CancerTools.org has made all reasonable efforts to ensure that the information provided by CancerTools.org and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.