

Product description

T47D-182R2 is an adherent human breast cancer cell line that exhibits resistance to fulvestrant (Faslodex). It was developed from the parental T47D/S5 cell line through long-term exposure to 100 nM fulvestrant. The cells retain an epithelial morphology. As resistance to fulvestrant—a common second-line endocrine therapy—is a clinical challenge, T47D-182R2 represents a valuable model for studying acquired resistance mechanisms. It serves as a robust tool for identifying targeted therapies against resistant tumor cells, as well as for evaluating strategies to prevent or delay resistance emergence.

Name: T47D-182R2 cell line

Organism: Human

Disease: Cancer

Cancer detailed: Breast cancer

Tissue: Breast

Parent cell line: T47D/S5

Growth properties: Adherent

Model: Cancer cell line

Donor: Female, Caucasian, 54Y

Production details: Human breast cancer cell line derived from T47D/S5 by long term treatment with 100 nM fulvestrant.

Cellosaurus ID: (CVCL_1D35)

Biosafety level: 1

Contributor(s)

Inventor: Anne Lykkesfeldt

Institute: Danish Cancer Society

Properties

Product format: Frozen

Unpacking and storage:

1. Check all containers for leakage or breakage.
2. Remove the frozen cells from the dry ice packaging and immediately place the cells at a temperature below -130°C, preferably in liquid nitrogen vapor, until ready for use.

Recommended medium: Phenol red free RPMI 1640 + 5% FCS + glutamax + 8ug Insulin/ml + 100 nM fulvestrant. Fetal Calf Serum (FCS) typically contains less estrogen than Fetal Bovine Serum (FBS) and is the preferred supplement for this cell line.

Culture conditions: 37.0°C ± 1.0°C humidified incubator with 5.0% CO₂

Cryopreservation medium: 10% DMSO in FCS

Handling instructions

1. Please ensure that vials are frozen when received, and store at **<-130 °C long term**. When removing frozen cells from storage, it is important to minimize exposure to room temperature (15 - 25°C). If not proceeding directly to thawing, place the cells on dry ice or in a liquid nitrogen container.
2. **Do not thaw at room temperature.** To thaw, swirl the vial quickly in a 37 °C water bath with O-ring and cap above the water to avoid contamination. Remove from the water bath with a small ice pellet remaining (this should not take more than 2 minutes) and wipe the exterior with 70% ethanol or isopropanol before transferring to a biosafety cabinet. Further steps should be conducted under aseptic conditions.
3. We strongly recommend that the volume of cell suspension is measured, and a 20 uL aliquot be set aside at this point for a viable cell count using trypan blue or similar dye.
9. Dilute the cell suspension with sufficient medium and distribute 5 mL each into T25 flasks to achieve a seeding density of 1.8 - 2.0 x 10⁴ / cm². Place in 37°C, 5% CO₂ incubator.
10. Change medium after 24 hours to remove residual DMSO and then every 2-3 days.
11. Subculture routine: Split 1:7 weekly (slow growing cell line) with Trypsin-EDTA for detachment at 37 °C for 5 minutes.

References

- Larsen et al. 2015. PLoS One. 10(2):e0118346. PMID: 25706943.
- Larsen et al. 2015. BMC Cancer. 15(1):1-15. PMID: 25885472.
- Kirkegaard et al. 2014. Cancer Lett. 344(1):90-100. PMID: 24513268.

Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: T47D-182R2 cell line, was invented by Anne Lykkesfeldt (CancerTools.org #151890).

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

While CancerTools.org has made all reasonable efforts to ensure that the information provided by CancerTools.org and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.